

**LOOSE HAMILTONIAN CYCLES FORCED BY LARGE $(k - 2)$ -DEGREE
– APPROXIMATE VERSION –**

JOSEFRAN DE OLIVEIRA BASTOS, GUILHERME OLIVEIRA MOTA, MATHIAS SCHACHT,
JAKOB SCHNITZER, AND FABIAN SCHULENBURG

ABSTRACT. We prove that for all $k \geq 3$ and $1 \leq \ell < k/2$, every k -uniform hypergraph \mathcal{H} on n vertices with $\delta_{k-2}(\mathcal{H}) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} + o(1)\right) \binom{n}{2}$ contains a Hamiltonian ℓ -cycle if $k - \ell$ divides n . This degree condition is asymptotically best possible.

§1. INTRODUCTION

A k -uniform hypergraph \mathcal{H} is a pair (V, E) with vertex set V and edge set E such that each edge is a subset of k vertices. Given a k -uniform hypergraph $\mathcal{H} = (V, E)$ and $S \in \binom{V}{s}$, we denote by $\deg(S)$ the number of edges of \mathcal{H} containing S and we denote by $N(S)$ the $(k - s)$ -element sets $T \in \binom{V}{k-s}$ such that $T \cup S$ is an edge of E , i.e. $\deg(S) = |N(S)|$. We define the *minimum s -degree* of \mathcal{H} , denoted by $\delta_s(\mathcal{H})$, as the minimum of $\deg(S)$ over all s -vertex sets $S \in \binom{V}{s}$.

We say that a k -uniform hypergraph \mathcal{C}_ℓ is an ℓ -cycle if there exists a cyclic ordering of its vertices such that every edge of \mathcal{C}_ℓ is composed of k consecutive vertices, two consecutive edges share exactly ℓ vertices, and every vertex is contained in an edge.

We are interested in the problem of finding minimum degree conditions that ensure the existence of Hamiltonian cycles. This problem was first studied by Katona and Kierstead in [6]. They posed a conjecture, which was confirmed by the following result of Rödl, Ruciński, and Szemerédi [11, 12]: For every $k \geq 3$, if \mathcal{H} is a k -uniform n -vertex hypergraph with $\delta_{k-1}(\mathcal{H}) \geq (1/2 + o(1))n$, then \mathcal{H} contains a Hamiltonian $(k - 1)$ -cycle. Their proof introduces the so-called *Absorbing Method*, which we will use in our proof as well. In [10] Kühn and Osthus investigated a similar question for 1-cycles, proving that 3-uniform hypergraphs \mathcal{H} with $\delta_2(\mathcal{H}) \geq (1/4 + o(1))n$ contain a Hamiltonian 1-cycle. This result

2010 *Mathematics Subject Classification.* 05C65 (primary), 05C45 (secondary).

Key words and phrases. hypergraphs, Hamiltonian cycles, degree conditions.

The first author was supported by CAPES. The second author was supported by FAPESP (Proc. 2013/11431-2 and 2013/20733-2) and CNPq (Proc. 477203/2012-4 and 456792/2014-7). The third author was supported through the Heisenberg-Programme of the DFG. The cooperation was supported by a joint CAPES/DAAD PROBRAL (Proc. 430/15).

was generalized to arbitrary k and ℓ -cycles with $1 \leq \ell < k/2$ by Hàn and Schacht [3] (see also [7]).

Theorem 1. *For all integers $k \geq 3$ and $1 \leq \ell < k/2$ and every $\gamma > 0$ there exists an n_0 such that every k -uniform hypergraph $\mathcal{H} = (V, E)$ on $|V| = n \geq n_0$ vertices with $n \in (k - \ell)\mathbb{N}$ and*

$$\delta_{k-1}(\mathcal{H}) \geq \left(\frac{1}{2(k-\ell)} + \gamma \right) n$$

contains a Hamiltonian ℓ -cycle. □

To see the optimality of the minimum degree condition, we consider the following well-known example. Let $\mathcal{H}_{k,\ell} = (V, E)$ be a k -uniform hypergraph on n vertices such that E is the set of all edges with at least one vertex from $A \subset V$, where $|A| = \left\lceil \frac{n}{2(k-\ell)} - 1 \right\rceil$. Note that an ℓ -cycle on n vertices contains $n/(k-\ell)$ edges and for $k < \ell/2$ every vertex is contained in at most two edges of any ℓ -cycle. So the hypergraph $\mathcal{H}_{k,\ell}$ does not contain a Hamiltonian ℓ -cycle and has $\delta_{k-1}(\mathcal{H}_{k,\ell}) = \left\lceil \frac{n}{2(k-\ell)} - 1 \right\rceil$. In [4] Han and Zhao proved a version of Theorem 1 with this sharp degree condition.

Kühn, Mycroft, and Osthus [9] generalized Theorem 1 to $1 \leq \ell < k - 1$, solving the problem of finding minimum $(k - 1)$ -degree conditions that ensure the existence of Hamiltonian ℓ -cycles in k -uniform hypergraphs. A natural question is to ask for minimum d -degree conditions forcing the existence of Hamiltonian ℓ -cycles for $d < k - 1$. In this direction Buß, Hàn, and Schacht proved the following asymptotically optimal result in [2].

Theorem 2. *For all $\gamma > 0$ there exists an n_0 such that every 3-uniform hypergraph $\mathcal{H} = (V, E)$ on $|V| = n \geq n_0$ vertices with $n \in 2\mathbb{N}$ and*

$$\delta_1(\mathcal{H}) \geq \left(\frac{7}{16} + \gamma \right) n$$

contains a Hamiltonian 1-cycle. □

Note that the optimality again follows from the hypergraph $\mathcal{H}_{k,\ell}$ considered above for $k = 3$ and $\ell = 1$. The sharp bound for $\delta_1(\mathcal{H})$ was proved by Han and Zhao in [5]. We generalize Theorem 2 to k -uniform hypergraphs and give an asymptotically optimal bound on the minimum $(k - 2)$ -degree for the existence of Hamiltonian ℓ -cycles for all $1 \leq \ell < k/2$.

Theorem 3 (Main result). *For all integers $k \geq 4$ and $1 \leq \ell < k/2$ and every $\gamma > 0$ there exists an n_0 such that every k -uniform hypergraph $\mathcal{H} = (V, E)$ on $|V| = n \geq n_0$ vertices with $n \in (k - \ell)\mathbb{N}$ and*

$$\delta_{k-2}(\mathcal{H}) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} + \gamma \right) \binom{n}{2}$$

contains a Hamiltonian ℓ -cycle.

The hypergraph $\mathcal{H}_{k,\ell}$ motivates the following notion of extremality. Let $k \geq 3$ and $\ell \geq 1$ be integers and let $0 < \xi < 1$. A k -uniform hypergraph $\mathcal{H} = (V, E)$ is called (ℓ, ξ) -extremal if there exists a set $B \subset V$ such that $|B| = \lfloor \frac{2(k-\ell)-1}{2(k-\ell)}n \rfloor$ and $e(B) \leq \xi \binom{n}{k}$, where $e(B)$ stands for the number of edges in the subhypergraph of \mathcal{H} induced by B . Our main result follows directly from the following theorem.

Theorem 4. *For any $0 < \xi < 1$ and all integers $k \geq 4$ and $1 \leq \ell < k/2$, there exists $\gamma > 0$ such that the following holds for sufficiently large n . Suppose \mathcal{H} is a k -uniform hypergraph on n vertices with $n \in (k - \ell)\mathbb{N}$ such that \mathcal{H} is not (ℓ, ξ) -extremal and*

$$\delta_{k-2}(\mathcal{H}) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} - \gamma \right) \binom{n}{2}.$$

Then \mathcal{H} contains a Hamiltonian ℓ -cycle.

We remark that for $k = 3$ and $\ell = 1$, the corresponding version of Theorem 4 appeared in the so-called non-extremal case of the sharp version of Theorem 2 in [4]. As a result, it will be sufficient to address the extremal case for a sharp version of Theorem 3 and we shall return to this in the near future [1]. For details about this approach see [4, 5]. It is easy to check that if $\delta_{k-2}(\mathcal{H}) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} + \gamma \right) \binom{n}{2}$, then there exists $\xi = \xi(k, \ell, \gamma) > 0$ such that \mathcal{H} is not (ℓ, ξ) -extremal. Consequently, Theorem 3 follows from Theorem 4.

§2. MAIN LEMMAS

2.1. Outline of the proof of Theorem 4. The proof follows the *Absorbing Method* introduced by Rödl, Ruciński, and Szemerédi in [11]. For this, we derive the following lemmas: the Absorbing Lemma (Lemma 7), the Reservoir Lemma (Lemma 6), and the Path-Tiling Lemma (Lemma 16).

We call an ℓ -path $\mathcal{A} \subseteq \mathcal{H}$ a β -absorbing path for a k -uniform hypergraph \mathcal{H} if for every subset $U \subset V(\mathcal{H})$ of size at most βn there exists an ℓ -path \mathcal{Q} such that $V(\mathcal{Q}) = V(\mathcal{A}) \cup U$ and \mathcal{Q} has the same ends as \mathcal{A} , for some $\beta > 0$. The Absorbing Lemma (Lemma 7) ensures the existence of a β -absorbing path \mathcal{A} . This reduces the problem of finding a Hamiltonian ℓ -cycle to that of finding an almost spanning ℓ -cycle that contains \mathcal{A} .

To obtain an almost spanning ℓ -cycle, we first find a bounded number (independent of $|V(\mathcal{H})|$) of ℓ -paths covering almost all vertices of $V(\mathcal{H}) \setminus \mathcal{A}$ and then connect them using only vertices from a small set, a so-called *reservoir set* that we fix beforehand. The Reservoir Lemma (Lemma 6) shows that it is possible to find this reservoir set R such that any bounded number of disjoint ℓ -paths can be connected to an ℓ -cycle, only using vertices from R .

We can choose the sizes of \mathcal{A} and R small enough, so that the remaining hypergraph satisfies almost the same degree condition as \mathcal{H} . Then the Path-Tiling Lemma (Lemma 16) ensures the existence of a collection of ℓ -paths covering almost all vertices of $V(\mathcal{H}) \setminus (\mathcal{A} \cup R)$. This is the only point in the proof where we use the exact value of the degree condition and the non-extremality of \mathcal{H} . (In fact, a proof for the corresponding version of the Path-Tiling Lemma for a direct proof of Theorem 3, which allows us to utilize a slightly larger degree condition, is a bit simpler.)

As mentioned before, the paths from the Path-Tiling Lemma and \mathcal{A} can be connected by using vertices from R to an almost spanning ℓ -cycle containing \mathcal{A} . Since this ℓ -cycle contains almost all vertices of \mathcal{H} , the absorbing property of \mathcal{A} allows us to absorb the leftover vertices, i.e. vertices that are not contained in any of the ℓ -paths and vertices that were not used to connect the ℓ -paths. The resulting ℓ -cycle is the desired Hamiltonian ℓ -cycle.

2.2. Connecting. In order to construct an almost spanning ℓ -cycle of a k -uniform hypergraph \mathcal{H} , we first find some ℓ -paths and connect them at their ends. Formally, given an ℓ -path $\mathcal{P} = v_1 \cdots v_t$ in \mathcal{H} , we define the *ends* of \mathcal{P} as the sets $\{v_1, \dots, v_\ell\}$ and $\{v_{t-\ell+1}, \dots, v_t\}$. For a collection of $2m$ mutually disjoint sets of ℓ vertices X_i, Y_i we say that a set of ℓ -paths $\mathcal{T}_1, \dots, \mathcal{T}_m$ *connects* $(X_i, Y_i)_{i \in [m]}$ if all paths are vertex-disjoint and X_i and Y_i are the ends of \mathcal{T}_i , for all $i \in [m]$. The connections for a given collection of disjoint ℓ -paths are given by the following lemma. In addition the lemma allows to restrict the edges used for the connection to a given “well-connected” subset R of vertices.

Lemma 5 (Connecting Lemma). *Let $\eta > 0$ and let $k \geq 4$, $1 \leq \ell < k/2$, and $m \geq 1$ be integers. Let $\mathcal{H} = (V, E)$ be a k -uniform hypergraph and $R \subset V$ with $|R| = r \geq 32km/\eta^3$. For every collection of $2m$ mutually disjoint sets $X_i, Y_i \in \binom{V}{\ell}$ and $V' = \bigcup_{i \in [m]} (X_i \cup Y_i) \cup R$ the following holds.*

If $|N(K) \cap \binom{R}{2}| \geq \eta \binom{r}{2}$ for all $K \in \binom{V'}{k-2}$, then there exist ℓ -paths $\mathcal{T}_1, \dots, \mathcal{T}_m$ of length at most 4 connecting $(X_i, Y_i)_{i \in [m]}$, which contain vertices from V' only.

Proof. Given $\eta > 0$ and integers $k \geq 4$, $1 \leq \ell < k/2$ and $m \geq 1$. Let $\mathcal{H} = (V, E)$, $R \subseteq V$, X_i, Y_i for $i \in [m]$, and V' satisfy the assumptions of the lemma.

Suppose we have constructed ℓ -paths $\mathcal{T}_1, \dots, \mathcal{T}_{j-1}$ connecting the pairs $(X_i, Y_i)_{i \in [j-1]}$ for some $j \leq m$ using only vertices from V' . We want to construct a path \mathcal{T}_j with ends X_j and Y_j . We define $F_j = \bigcup_{i \in [m]} (X_i \cup Y_i) \cup \bigcup_{i \in [j-1]} V(\mathcal{T}_i)$ as the set of *forbidden vertices* for \mathcal{T}_j .

If $k-2 \geq 2\ell = |X_j \cup Y_j|$, fix a set Z of size $k-2-2\ell$ from $R \setminus F_j$. Since $|R| = r \geq 32km/\eta^3$, we know that

$$\left| N(X_j \cup Y_j \cup Z) \cap \binom{R}{2} \right| \geq \eta \binom{r}{2} > \binom{r}{2} - \binom{r-4km}{2} \geq \binom{r}{2} - \binom{|R \setminus F_j|}{2}.$$

Hence, there exists a hyperedge $X_j \cup Y_j \cup Z'$ with $Z' \subseteq R \setminus F_j$, which realizes the path \mathcal{T}_j .

It is left to consider the case that $2\ell = k-1$. See Figure 1 for a drawing of the path we will construct in this case. For a set $A \subseteq V$, let $N_A(S) = N(S) \cap \binom{A}{k-s}$.

Observation. *For any $Z \in \{X_j, Y_j\}$ and $L \in \binom{R \setminus F_j}{\ell-1}$, there are at least $\eta r/4$ many vertices $z \in R \setminus (F_j \cup L)$ with $|N_{R \setminus F_j}(Z \cup L \cup \{z\})| \geq \eta r/4$.*

To see the observation note that we can consider $N_{R \setminus F_j}(Z \cup L)$ as the edge set of a 2-graph with vertex set $R \setminus (F_j \cup L)$. Since $r \geq 32km/\eta^3$, it follows from the degree condition of \mathcal{H} into the set R that this graph has edge density at least $\eta/2$ and the observation follows.

Let $L \in \binom{R \setminus F_j}{\ell-1}$ and let $x, y \in R \setminus F_j$ be distinct. We say that (x, L, y) is an *extendable triple* in $R \setminus F_j$ if

$$|N_{R \setminus F_j}(X_j \cup L \cup \{x\})| \geq \eta r/4 \quad \text{and} \quad |N_{R \setminus F_j}(Y_j \cup L \cup \{y\})| \geq \eta r/4.$$

The observation yields at least $(\eta r/4)(\eta r/4 - 1) > (\eta r/8)^2$ extendable triples (x, L, y) for any fixed $L \in \binom{R \setminus F_j}{\ell-1}$.


Given $S \in \binom{R \setminus F_j}{\ell-2}$ and an extendable triple (x, L, y) disjoint from S , i.e. $S \cup L \cup \{x, y\}$ is a $(k-2)$ -element set. Consequently, the minimum degree condition of the lemma yields at least $\eta \binom{r}{2}$ pairs $M \in \binom{R}{2}$ such that $S \cup M \cup L \cup \{x, y\}$ is an edge of \mathcal{H} . Moreover, similar as in the proof of the observation at least $(\eta/2) \binom{|R \setminus F_j|}{2}$ of these pairs avoid F_j . Since this is true for every extendable triple and there are at least $\binom{|R \setminus F_j|}{\ell-1} (\eta r/8)^2$ extendable triples, there exists an $M \in \binom{R \setminus F_j}{2}$ that forms an edge of \mathcal{H} together with at least $(\eta/2)(\eta r/8)^2 \binom{|R \setminus F_j|}{\ell-1}$ extendable triples. Since $r \geq 32km/\eta^3$, this is more than the number of triples that any single extendable triple can intersect with, so there exist two completely disjoint extendable triples (x, L, y) and (x', L', y') that form an edge of \mathcal{H} together with $M' = M \cup S$.

By the definition of extendable triples we have

$$|N_{R \setminus F_j}(X_j \cup L \cup \{x\})| \geq \eta r/4 > k+1 = |M' \cup L' \cup \{x', y', y\}|$$

and

$$|N_{R \setminus F_j}(Y_j \cup L' \cup \{y'\})| \geq \eta r/4 > k+2 = |M' \cup L \cup \{x, y, x'\}| + 1.$$

FIGURE 1. The path connecting (X_j, Y_j) .

Consequently there are $v, v' \in R \setminus F_j$ such that the hyperedges

$$\{X_j \cup L \cup \{v, x\}\}, \quad \{M' \cup L \cup \{x, y\}\}, \quad \{M' \cup L' \cup \{x', y'\}\}, \quad \text{and} \quad \{Y_j \cup L' \cup \{y', v'\}\}$$

are edges of \mathcal{H} , which form a path of length 4 connecting (X_j, Y_j) . \square

In the main proof we will connect ℓ -paths to an almost spanning ℓ -cycle. The Reservoir Lemma (stated below) ensures the existence of a small set R such that we can connect an arbitrary collection of at most $2m$ many ℓ -sets, only using vertices of R .

Lemma 6 (Reservoir Lemma). *Let $\eta, \varepsilon > 0$ and let $k \geq 4$, $1 \leq \ell < k/2$, and $m \geq 1$ be integers. Then for every sufficiently large k -uniform hypergraph $\mathcal{H} = (V, E)$ on n vertices with $\delta_{k-2}(\mathcal{H}) \geq \eta \binom{n}{2}$ there is a set $R \subset V$ with $|R| \leq \varepsilon n$ such that the following holds.*

For every collection X_i, Y_i for $i \in [j]$ of $2j$ mutually disjoint sets of ℓ vertices, where $j \leq m$, there exist ℓ -paths $\mathcal{T}_1, \dots, \mathcal{T}_j$ of length at most 4 connecting $(X_i, Y_i)_{i \in [j]}$ that, moreover, contain vertices from $\bigcup_{i \in [j]} (X_i \cup Y_i) \cup R$ only.

Lemma 6 is a consequence of Lemma 5, since one can show that a suitably sized random subset $R \subseteq V$ inherits with high probability an appropriately scaled minimum degree condition from \mathcal{H} . As a consequence such a set satisfies the assumptions of Lemma 5 and the lemma yields the conclusion of Lemma 6 (see, e.g. [2, Lemma 6] for a very similar argument).

2.3. Absorption. Given a k -uniform hypergraph \mathcal{H} and $U \subset V$ with $|U| \in (k - \ell)\mathbb{N}$, we say that an ℓ -path \mathcal{A} *absorbs* U if there exists an ℓ -path \mathcal{Q} with the same ends as \mathcal{A} and $V(\mathcal{Q}) = V(\mathcal{A}) \cup U$. At the end of the main proof we will absorb all vertices outside of an almost spanning ℓ -cycle to obtain a Hamiltonian ℓ -cycle using an *absorbing path* \mathcal{A} , i.e. a path that can absorb any set U of small linear size. The existence of such a path \mathcal{A} is given by the following lemma.

Lemma 7 (Absorbing Lemma). *For every $\eta, \zeta > 0$ and all integers $k \geq 4$ and $1 \leq \ell < k/2$ there exists $\varepsilon > 0$ such that the following holds for sufficiently large n . Let $\mathcal{H} = (V, E)$ be a k -uniform hypergraph on n vertices that satisfies $\delta_{k-2}(\mathcal{H}) \geq \eta \binom{n}{2}$. Then there is an ℓ -path \mathcal{A} with $|V(\mathcal{A})| \leq \zeta n$ such that for all subsets $U \subset V \setminus V(\mathcal{A})$ of size at most εn with $|U| \in (k-\ell)\mathbb{N}$ there exists an ℓ -path $\mathcal{Q} \subset \mathcal{H}$ with $V(\mathcal{Q}) = V(\mathcal{A}) \cup U$ such that \mathcal{A} and \mathcal{Q} have the same ends.*

Proof. Let $\eta, \zeta > 0$ and let $k \geq 4$ and $1 \leq \ell < k/2$ be integers. Fix auxiliary constants

$$\tilde{\eta} = \frac{\eta}{4k!} \quad \text{and} \quad t = 3k - 2\ell$$

and set

$$\varepsilon = \frac{\zeta \tilde{\eta}^{10}}{56k}.$$

Let n be sufficiently large and let $\mathcal{H} = (V, E)$ be a k -uniform hypergraph on n vertices that satisfies $\delta_{k-2}(\mathcal{H}) \geq \eta \binom{n}{2}$. First, we will show that for any $S \in \binom{V}{k-\ell}$ there exist many, i.e. $\Omega(n^t)$, 3-edge ℓ -paths that absorb S (see Claim 8 below). For that we will use the following consequence of the minimum degree condition. Let $A, B \subset V(\mathcal{H})$ be disjoint sets of vertices with $|A| \leq k-2$ and $|B| \leq t+k$. Then,

$$\deg_{\mathcal{H}[V \setminus B]}(A) \geq \frac{(n-|A|) \cdot \dots \cdot (n-k+3)}{(k-|A|)!} \cdot \eta \binom{n}{2} - |B|n^{k-|A|-1} \geq \tilde{\eta} n^{k-|A|}. \quad (1)$$

Claim 8. *For every $S \in \binom{V}{k-\ell}$ there exist at least $\tilde{\eta}^5 n^t$ many 3-edge ℓ -paths that absorb S .*

Proof. Let $S_1 \cup S_2 = S$ chosen in some way such that

$$|S_1| \geq |S_2| \geq |S_1| - 1 \quad \text{and} \quad \max\{0, 3\ell - k\} \leq |S_1 \cap S_2| < \ell \quad (2)$$

and set $s_1 = |S_1|$, $s_2 = |S_2|$, and $s_3 = |S_1 \cap S_2|$. Clearly, we have

$$s_1 + s_2 - s_3 = |S| = k - \ell. \quad (3)$$

It follows from the choices above that $s_1 + s_2 \geq 2\ell$. Indeed, since $s_3 \geq 3\ell - k$ we have $k - \ell = s_1 + s_2 - s_3 \leq s_1 + s_2 - 3\ell + k$ and, hence, $s_1 + s_2 \geq 2\ell$. Furthermore, $s_1 \geq s_2 \geq s_1 - 1$ yields

$$s_1 \geq s_2 \geq \ell. \quad (4)$$

We then select the following sets. See Figure 2 for a drawing of the chosen sets and edges containing them. In each step, we will only select sets that are disjoint from S and anything chosen in a previous step.




FIGURE 2. The path \mathcal{P} , consisting of e_1, e_2 , and e_3 , that absorbs S .

- (i) Since $s_1 \leq k - \ell - 1 \leq k - 2$, by (1) there exist $\tilde{\eta}n^{k-s_1}$ choices for a $(k - s_1)$ -set X such that $f_1 = X \cup S_1$ is an edge of \mathcal{H} . Since $|X| = k - s_1 \stackrel{(3)}{=} \ell + s_2 - s_3$ it follows from (4) that we may partition $X = L_1 \cup F \cup F_1$ such that $|L_1| = \ell$ and $|F| = \ell - s_3$.
- (ii) Since $k \geq 4$ we have $k - \ell \geq 3$ and, consequently, $s_1 \geq \lceil (k - \ell)/2 \rceil \geq 2$. Thus, by (1) and $|S_2 \cup F| = s_2 + \ell - s_3 = k - s_1$, there exist $\tilde{\eta}n^{s_1}$ choices for a set Y of size s_1 such that $f_2 = S_2 \cup F \cup Y$ is an edge of \mathcal{H} . Again owing to (4) we may partition $Y = L_2 \cup F_2$ such that $|L_2| = \ell$.
- (iii) Fix $L'_1 \subset L_1$ and $L'_2 \subset L_2$ subsets of size $\ell - 1$. Note that

$$|L'_1 \cup L'_2 \cup F \cup F_1 \cup F_2| = |X| + |Y| - 2 = k - 2.$$

Therefore, there exist at least $\tilde{\eta}n^2$ choices for a pair of vertices $\{x_1, x_2\}$ such that $e_2 = \{x_1, x_2\} \cup L'_1 \cup L'_2 \cup F \cup F_1 \cup F_2$ is an edge of \mathcal{H} .

- (iv) Since $k \geq 4$ we have $\ell + 1 \leq k - 2$. Therefore, there exist $\tilde{\eta}n^{k-(\ell+1)}$ choices each for two disjoint edges e_1 and e_3 such that $\{x_1\} \cup L_1 \subset e_1$ and $\{x_2\} \cup L_2 \subset e_3$.

By construction we have

$$e_1 \cap e_2 = \{x_1\} \cup L'_1 \quad \text{and} \quad e_2 \cap e_3 = \{x_2\} \cup L'_2,$$

so the edges e_1, e_2 , and e_3 form an ℓ -path \mathcal{P} in \mathcal{H} . Moreover, since

$$e_1 \cap f_1 = L_1, \quad |f_1 \cap f_2| = |(S_1 \cap S_2) \cup F| \stackrel{(i)}{=} \ell, \quad \text{and} \quad f_2 \cap e_3 = L_2,$$

the edges e_1, f_1, f_2 , and e_3 form an ℓ -path \mathcal{P}' . Since $k - \ell - 1 \geq \ell$, we may select for \mathcal{P} and \mathcal{P}' the same ends in e_1 and e_3 . Moreover, $V(\mathcal{P}') = V(\mathcal{P}) \cup S$ and, therefore, the ℓ -path \mathcal{P} absorbs S . From (i)–(iv) it is clear that there are at least $\tilde{\eta}^5 n^t$ choices for \mathcal{P} . \square

Following the scheme from [11], let $\mathcal{F} \subset V(\mathcal{H})^t$ be a family of ordered t -sets of vertices such that each of these sets are selected from $V(\mathcal{H})^t$ independently with probability

$$p = \frac{4\varepsilon}{\tilde{\eta}^5 n^{t-1}}.$$

An ℓ -path in $V(\mathcal{H})^t$ is an ordered set (v_1, \dots, v_t) of vertices such that

$$e_1 = \{v_1, \dots, v_k\}, \quad e_2 = \{v_{k-\ell+1}, \dots, v_{2k-\ell}\}, \quad \text{and} \quad e_3 = \{v_{2k-2\ell+1}, \dots, v_{3k-2\ell}\}$$

are edges in \mathcal{H} . Using Chernoff's inequality, with high probability we have

$$|\mathcal{F}| \leq 2pn^t.$$

By Claim 8, for each set S of size $k - \ell$, at least $\tilde{\eta}^5 n^t$ ℓ -paths in $V(\mathcal{H})^t$ absorb S . By Chernoff's inequality, w.h.p. for all $S \in \binom{V}{k-\ell}$, there are at least $2\varepsilon n$ ℓ -paths in \mathcal{F} that absorb S . Furthermore, by Markov's inequality, the number of intersecting pairs of t -sets in \mathcal{F} is at most εn with probability at least $1/2$.

Let \mathcal{F} be a family that satisfies the above conditions. For each of the intersecting pairs in \mathcal{F} , delete one of the t -sets and let $\mathcal{F}' \subset \mathcal{F}$ be the remaining family. Using Lemma 5 (with $R = V$) we can connect all ℓ -paths in \mathcal{F}' to an ℓ -path \mathcal{A} with

$$|V(\mathcal{A})| \leq |\mathcal{F}'| \cdot (4k + t) \leq 2pn^t \cdot 7k = \frac{56k}{\tilde{\eta}^5} \varepsilon n \leq \zeta n$$

and this path absorbs all sets $U \subset V \setminus V(\mathcal{A})$ with $|U| \in (k - \ell)\mathbb{N}$ and $|U| \leq \varepsilon n$. \square

2.4. Path-Tiling. In this part we will find a path-tiling of ℓ -paths in \mathcal{H} that covers all but a small fraction of the vertices of \mathcal{H} . For that purpose we use the so-called *weak regularity lemma* for hypergraphs, which is the straightforward extension of Szemerédi's regularity lemma for graphs [13]. Roughly speaking, we will show that there exists a fractional \mathcal{C}_ℓ -tiling, a so-called β -hom(\mathcal{C}_ℓ)-tiling in the resulting reduced hypergraph \mathcal{R} of \mathcal{H} , where \mathcal{C}_ℓ is the k -uniform "cherry" consisting of two hyperedges that share exactly 2ℓ vertices. The fractional \mathcal{C}_ℓ -tiling of \mathcal{R} will transfer to a path-tiling of \mathcal{H} .

First, we introduce the standard notation for the regularity lemma. Let $\mathcal{H} = (V, E)$ be a k -uniform hypergraph and let V_1, \dots, V_k be non-empty, mutually disjoint subsets of V . We denote the number of edges with one vertex in each V_i by $e_{\mathcal{H}}(V_1, \dots, V_k)$ and define the density of \mathcal{H} w.r.t. (V_1, \dots, V_k) by

$$d_{\mathcal{H}}(V_1, \dots, V_k) = \frac{e_{\mathcal{H}}(V_1, \dots, V_k)}{|V_1| \cdots |V_k|}.$$

For $\varepsilon > 0$ and $d > 0$, a k -tuple (V_1, \dots, V_k) of mutually disjoint subsets of vertices is called (ε, d) -regular if for all k -tuples (A_1, \dots, A_k) of subsets $A_i \subseteq V_i$ with $|A_i| \geq \varepsilon |V_i|$, we have

$$|d_{\mathcal{H}}(A_1, \dots, A_k) - d| \leq \varepsilon.$$

Moreover, the tuple (V_1, \dots, V_k) is called ε -regular if it is (ε, d) -regular for some $d > 0$.

Lemma 9 (Weak regularity lemma). *For all integers $k \geq 2$ and $t_0 \geq 1$ and for every $\varepsilon > 0$, there exists $T_0 = T_0(k, t_0, \varepsilon)$ such that for every sufficiently large k -uniform hypergraph $\mathcal{H} = (V, E)$ on n vertices, there exists a partition $V = V_0 \cup V_1 \cup \dots \cup V_t$ satisfying*

- (i) $t_0 \leq t \leq T_0$,
- (ii) $|V_1| = \dots = |V_t|$ and $|V_0| \leq \varepsilon n$, and
- (iii) for all but at most $\varepsilon \binom{t}{k}$ many k -subsets $\{i_1, \dots, i_k\} \subset [t]$, the k -tuple $(V_{i_1}, \dots, V_{i_k})$ is ε -regular.

A vertex partition of a hypergraph \mathcal{H} satisfying (i)–(iii) of the conclusion of Lemma 9 will be referred to as an ε -regular partition. For $\varepsilon > 0$ and $d > 0$, we define the *reduced hypergraph* $\mathcal{R} = \mathcal{R}(\varepsilon, d)$ of \mathcal{H} w.r.t. such a partition as the k -uniform hypergraph on the vertex set $[t]$ and

$$\{i_1, \dots, i_k\} \in E(\mathcal{R}) \iff (V_{i_1}, \dots, V_{i_k}) \text{ is } (\varepsilon, d')\text{-regular, for some } d' \geq d.$$

In typical applications of the regularity lemma, the reduced hypergraph inherits some key features of the given hypergraph \mathcal{H} . In fact, the following observation shows that the reduced hypergraph inherits approximately the minimum degree condition of the original hypergraph. A similar result can be found in [3, Proposition 16] and for completeness we include its proof below.

Lemma 10. *Given $c, \varepsilon, d > 0$ and integers $k \geq 3$ and $t_0 \geq 2k/d$. Let \mathcal{H} be a k -uniform hypergraph on $n \geq t \geq t_0$ vertices such that*

$$\delta_{k-2}(\mathcal{H}) \geq c \binom{n}{2}.$$

If \mathcal{H} has an ε -regular partition $V_0 \cup V_1 \cup \dots \cup V_t$ with reduced hypergraph $\mathcal{R} = \mathcal{R}(\varepsilon, d)$, then at most $\sqrt{\varepsilon} \binom{t}{k-2}$ many $(k-2)$ -subsets K of $[t]$ violate

$$\deg_{\mathcal{R}}(K) \geq (c - 2d - \sqrt{\varepsilon}) \binom{t}{2}.$$

Proof. Let $\mathcal{D} = \mathcal{D}(d)$ and $\mathcal{N} = \mathcal{N}(\varepsilon)$ be the hypergraphs with vertex set $[t]$ and

- $E(\mathcal{D})$ consists of all sets $\{i_1, \dots, i_k\}$ such that $d(V_{i_1}, \dots, V_{i_k}) \geq d$,
- $E(\mathcal{N})$ consists of all sets $\{i_1, \dots, i_k\}$ such that $(V_{i_1}, \dots, V_{i_k})$ is not ε -regular.

Note that the reduced hypergraph $\mathcal{R}(\varepsilon, d)$ is the hypergraph with vertex set $[t]$ and edge set $E(\mathcal{D}) \setminus E(\mathcal{N})$. For an arbitrary $K = \{i_1, \dots, i_{k-2}\} \in \binom{[t]}{k-2}$ we will show that

$$\deg_{\mathcal{D}}(K) \geq (c - 2d) \binom{t}{2}. \tag{5}$$

Let $n/t \geq |V_{i_j}| = m \geq (1-\varepsilon)n/t$ be the size of the partition classes and let x be the number of edges in \mathcal{H} that intersect each V_{i_j} exactly in one vertex for $j \in [k-2]$. By the condition on $\delta_{k-2}(\mathcal{H})$ and $t \geq t_0 \geq 2k/d$, we obtain

$$x \geq m^{k-2} \left(c \binom{n}{2} - (k-2)mn \right) \geq (c-d) m^{k-2} \binom{n}{2}.$$

If (5) would not hold, then we find for x the upper bound

$$x < (c-2d) \binom{t}{2} m^k + \binom{t}{2} dm^k \leq (c-d) m^{k-2} \binom{n}{2}$$

contradicting the lower bound for x .

Next we observe that at most $\sqrt{\varepsilon} \binom{t}{k-2}$ many $(k-2)$ -sets K satisfy $\deg_{\mathcal{N}}(K) \leq \sqrt{\varepsilon} \binom{t}{2}$ since the number of non- ε -regular k -tuples in \mathcal{R} is at most $\varepsilon \binom{t}{k}$. Consequently, it follows from the degree conditions on \mathcal{D} and \mathcal{N} that all but at most $\sqrt{\varepsilon} \binom{t}{k-2}$ many $(k-2)$ -sets K satisfy

$$\deg_{\mathcal{R}}(K) \geq (c-2d-\sqrt{\varepsilon}) \binom{t}{2}.$$

□

We will find a suitable fractional \mathcal{C}_ℓ -tiling in the reduced hypergraph \mathcal{R} , where the cherry \mathcal{C}_ℓ is the k -uniform hypergraph with vertex set $[2k-2\ell]$ and edges $\{1, \dots, k\}$ and $\{k-2\ell+1, \dots, 2k-2\ell\}$.

Definition 11. Let \mathcal{C} and \mathcal{R} be k -uniform hypergraphs, $\beta > 0$, and let Φ be a multiset of hypergraph homomorphisms from \mathcal{C} to \mathcal{R} . A function $h: \Phi \rightarrow \{a\beta: a \in \mathbb{N}_{>0}\}$ is called a β -hom(\mathcal{C})-tiling if

$$w_h(v) = \sum_{u \in V(\mathcal{C})} \sum_{\varphi \in \Phi: v = \varphi(u)} h(\varphi) \leq 1$$

for all $v \in V(\mathcal{R})$. We call

$$w(h) = \sum_{v \in V(\mathcal{R})} w_h(v) = \sum_{\varphi \in \Phi} h(\varphi) |V(\mathcal{C})|$$

the weight of the tiling.

The following building block allows us to easily define a tiling on a single edge.

Fact 12. Given an edge $e = \{v_1, \dots, v_k\}$, there exists a $\frac{1}{2(k-\ell-1)}$ -hom(\mathcal{C}_ℓ)-tiling h that is non-zero only on e , such that $w_h(v_i) = 1$ for $i \in [k-2]$ and $w_h(v_{k-1}) = w_h(v_k) = \frac{k-2}{2(k-\ell-1)}$. Note that we may scale the weight of h by any $q \in (0, 1]$ and obtain a $\frac{q}{2(k-\ell-1)}$ -hom(\mathcal{C}_ℓ)-tiling with $w_h(v_i) = q$ for $i \in [k-2]$ and $w_h(v_{k-1}) = w_h(v_k) = \frac{q(k-2)}{2(k-\ell-1)}$. Similarly, for any $q \in (0, 1]$ there exists a $\frac{q}{2(k-\ell)}$ -hom(\mathcal{C}_ℓ)-tiling with $w_h(v_i) = q$ for $i \in [k]$.

Proof. For this consider the homomorphism that maps \mathcal{C}_ℓ to e such that $v_1, \dots, v_{2\ell-2}, v_{k-1}$ and v_k are the image of the intersection of the two edges of \mathcal{C}_ℓ . By cyclically shifting the image of the first $2\ell - 2$ vertices of the intersection and appropriate scaling, we obtain all homomorphisms for the required tiling. We obtain the even weight distribution for the last part of the fact by cyclically shifting the whole image k times. \square

The following lemma is the main part of the proof of the Path-Tiling Lemma. For this we introduce a fractional notion of extremality. We say that a k -uniform hypergraph \mathcal{R} on t vertices is β -fractionally (ℓ, ξ) -extremal if there is a function $b: V(\mathcal{R}) \rightarrow \{0\} \cup [\beta, 1]$ with

$$\sum_{v \in V(\mathcal{R})} b(v) \geq \frac{2(k-\ell)-1}{2(k-\ell)}t \quad \text{and} \quad \sum_{e \in E(\mathcal{R})} \prod_{v \in e} b(v) \leq \xi \binom{t}{k}.$$

Note that the function b can be viewed as a set of weighted vertices, which plays the rôle of the vertex set B in the definition of extremality.

Lemma 13. *For all integers $k \geq 3$ and $1 \leq \ell < k/2$, there exist C and γ_0 such that for all $\alpha > 0$ and $\gamma \in (0, \gamma_0)$, there exist $\beta > 0$ and $\varepsilon > 0$ such that the following holds for sufficiently large t . Let \mathcal{R} be a k -uniform hypergraph on t vertices that is not β -fractionally $(\ell, C\gamma)$ -extremal and*

$$\deg(K) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} - \gamma \right) \binom{t}{2} \quad (6)$$

holds for all but at most $\varepsilon \binom{t}{k-2}$ sets $K \in \binom{V(\mathcal{R})}{k-2}$. Then there exists a β -hom(\mathcal{C}_ℓ)-tiling h with weight at least $(1-\alpha)t$.

Proof. Clearly, it is sufficient to prove the lemma for small values of α . Consequently the quantification of the lemma allows us to fix the parameters and auxiliary constants C' and c to satisfy the following hierarchy of constants

$$\frac{1}{k}, \frac{1}{\ell} \gg \frac{1}{C'} \gg \frac{1}{C} \gg \gamma_0 \geq \gamma \gg \alpha \gg c, \varepsilon. \quad (7)$$

Moreover, we fix β inductively such that

$$1 = \beta_0 \gg \beta_1 \gg \dots \gg \beta_{\lfloor 1/c \rfloor} = \beta \quad \text{and} \quad \frac{\beta_i}{\beta_{i+1}} \in \mathbb{N}$$

and let t be sufficiently large such that $c, \beta \gg 1/t$. Note that any β_i -hom(\mathcal{C}_ℓ)-tiling is also a β -hom(\mathcal{C}_ℓ)-tiling as β_i is a multiple of β . To prove the lemma, we show that given a β_i -hom(\mathcal{C}_ℓ)-tiling h with weight $w(h) < (1-\alpha)t$, there exists a β_{i+1} -hom(\mathcal{C}_ℓ)-tiling h' with weight $w(h') \geq w(h) + ct$. We can begin with the trivial 1-hom(\mathcal{C}_ℓ)-tiling with weight zero and hence, after at most $\frac{1}{c}$ steps, we obtain a β -hom(\mathcal{C}_ℓ)-tiling with weight at least $(1-\alpha)t$.

For the rest of the proof fix a β_i -hom(\mathcal{C}_ℓ)-tiling h with weight $w(h) < (1-\alpha)t$ and assume for a contradiction that there is no β_{i+1} -hom(\mathcal{C}_ℓ)-tiling with weight $w(h) + ct$. It

follows from the upper bound on the weight that there are at least $\alpha t/2$ vertices $v \in V(\mathcal{R})$ with $w_h(v) < 1 - \alpha/2$ and we may fix a subset W of them of size $\alpha t/2$. Although a tiling with bigger weight implies the existence of more edges, we will not use these edges for an improvement. So we may actually assume w.l.o.g. that

$$(1 - 2\alpha)t \leq w(h) < (1 - \alpha)t,$$

as we can otherwise add edges on $V \setminus W$ until this weight can be trivially achieved and remove these edges after the improvement step.

We view Φ , the set of homomorphisms from \mathcal{C}_ℓ to \mathcal{R} , as a multiset, where we include φ with multiplicity $\frac{h(\varphi)}{\beta_i}$, so that we can assume $h: \Phi \rightarrow \{\beta_i\}$. With this notion the following bounds on the size of Φ follow from the above

$$(1 - 2\alpha)\frac{t}{\beta_i v(\mathcal{C}_\ell)} \leq |\Phi| < (1 - \alpha)\frac{t}{\beta_i v(\mathcal{C}_\ell)}. \quad (8)$$

Also, we identify a homomorphism φ in Φ with the – not necessarily distinct – vertices $(v_1, \dots, v_{2k-2\ell})$ in its image, where $\{v_1, \dots, v_k\}$ and $\{v_{k-2\ell+1}, \dots, v_{2k-2\ell}\}$ form edges in \mathcal{R} . We refer to the elements of Φ as *cherries* $\mathcal{C} \in \Phi$.

Consider the $(k - 2)$ -sets in W that satisfy the degree condition (6) of the lemma. Since $\varepsilon \ll \alpha$, among those $(k - 2)$ -sets we find a collection \mathcal{W} whose elements are pairwise disjoint and cover at least $|W|/2$ vertices. For later reference we note

$$|\mathcal{W}| \geq \frac{|W|}{2(k - 2)} > \frac{\alpha t}{4k}. \quad (9)$$

For $K \in \mathcal{W}$ we consider the *link graph* L_K of K in \mathcal{R} , which is the (2-uniform) graph containing all edges e such that $K \cup e \in E(\mathcal{R})$. We will find pairs $\mathcal{C}, \mathcal{C}' \in \Phi$ on which the link graph allows us to find a better tiling. For this we only want to consider edges in the bipartite induced link graph $L_K(\mathcal{C}, \mathcal{C}')$, i.e. edges that have one vertex in \mathcal{C} and one in \mathcal{C}' . At most $\frac{t}{\beta_i} \binom{v(\mathcal{C}_\ell)}{2} \leq \gamma \binom{t}{2}$ edges have both ends in the same $\mathcal{C} \in \Phi$ and at most $\alpha t^2/2 \leq \gamma \binom{t}{2}$ edges contain a vertex from W , so let L'_K be the graph obtained from L_K by removing all these edges. Combined with the degree condition (6) we have

$$e(L'_K) \geq \left(\frac{4(k - \ell) - 1}{4(k - \ell)^2} - 3\gamma \right) \binom{t}{2} \quad (10)$$

for every such $(k - 2)$ -set $K \in \mathcal{W}$.

We show in the following that for most $K \in \mathcal{W}$ the bipartite link graph between most \mathcal{C} and \mathcal{C}' has a very specific structure. We call $\mathcal{C}, \mathcal{C}' \in \Phi$ an *extremal pair for K* if there exist special vertices $u \in \mathcal{C}$ and $u' \in \mathcal{C}'$ such that $L_K(\mathcal{C}, \mathcal{C}')$ contains exactly all edges incident to these two vertices.

Claim 14. *There exists an element $u_C \in \mathcal{C}$ for all $C \in \Phi$ such that the following holds. For all but $\gamma|\mathcal{W}|$ sets $K \in \mathcal{W}$ all but a $C'\gamma$ fraction of the pairs $C, C' \in \Phi$ are extremal for K with special vertices u_C and $u_{C'}$.*

Proof. The proof of the claim consists of three parts. First we show that if for a given $(k-2)$ -tuple $K \in \mathcal{W}$ and some pairs of cherries $C, C' \in \Phi$ the induced bipartite link graph $L_K(C, C')$ contains more than $4(k-\ell) - 1$ edges, then there is a *local improvement* of the tiling by a weight of at least $\beta_i/4$. In a second step we shall bound the number of possible local improvements, as otherwise we could combine them to arrive at a desired tiling h' with a weight increased by ct and this would already conclude the proof of Lemma 13. In the last step we utilise this bound on the number of local improvements to show that “typically” $L_K(C, C')$ contains only $4(k-\ell) - 1$ edges and displays the structural conditions stated in the claim.

For the first step we consider two cases. Suppose that there is a matching with three edges in $L_K(C, C')$ for some $K \in \mathcal{W}$ and $C, C' \in \Phi$. Recall that $L_K(C, C')$ is a bipartite graph with partition classes of size $2(k-\ell)$. Then we assign the weight $\frac{k-2}{6(k-\ell-1)}\beta_i$ to the vertices of the matching edges and the weight β_i to the vertices of K . Setting in addition $h'(C) = h'(C') = (1 - \frac{k-2}{6(k-\ell-1)})\beta_i$, this defines a valid β_{i+1} -hom-tiling h' by applying Fact 12 (with $q = \frac{1}{3}\beta_i$) to the three edges in \mathcal{R} corresponding to the matching edges in the link graph. Indeed, by considering the vertices on which the weight has changed, it is easy to see that

$$w(h') = w(h) + \left(k - 2 - (4k - 4\ell - 6) \cdot \frac{k-2}{6(k-\ell-1)} \right) \beta_i \geq w(h) + \frac{1}{3}\beta_i,$$

which yields the local improvement in this case.

In the other case, suppose that there is no matching of size three in $L_K(C, C')$. Then by König’s theorem [8] there is a vertex cover of size at most two. Consequently, there are at most $4(k-\ell)$ edges in $L_K(C, C')$. Note that if $L_K(C, C')$ contains indeed $4(k-\ell)$ edges, then the two cover vertices are in the same class – say in C – and their neighbourhood is all of C' . In particular there are two edges for each of the two vertices using four distinct vertices in C' . On the vertices of these edges we put weights $\frac{k-2}{8(k-\ell-1)}\beta_i$ and β_i on the vertices of K . Set $h'(C) = (1 - \frac{k-2}{4(k-\ell-1)})\beta_i$ and $h'(C') = (1 - \frac{k-2}{8(k-\ell-1)})\beta_i$. Again, this defines a tiling h' with

$$\begin{aligned} w(h') &= w(h) + \left(k - 2 - (2k - 2\ell - 2) \cdot \frac{k-2}{4(k-\ell-1)} - (2k - 2\ell - 4) \cdot \frac{k-2}{8(k-\ell-1)} \right) \beta_i \\ &\geq w(h) + \frac{k-2}{4}\beta_i \geq w(h) + \frac{1}{4}\beta_i. \end{aligned}$$

This establishes the local improvement for this case and concludes the first step.

For the second step suppose that there are at least $\gamma|\mathcal{W}|/2$ many $K \in \mathcal{W}$ such that for each such K we can define a local improvement for $\gamma|\Phi|^2$ cherry pairs. We apply these local improvements greedily, only using each cherry $\mathcal{C} \in \Phi$ at most once (over all K), to increase the weight of the tiling as much as possible. This procedure may end, either when every $K \in \mathcal{W}$ contains a saturated vertex and in this case we enlarge the total weight by at least $\frac{\alpha}{2} \cdot \gamma|\mathcal{W}|/2$, or when for every $K \in \mathcal{W}$ for each of the $\gamma|\Phi|^2$ pairs of cherries at least one cherry was used for some local improvement. Since any collection of $\gamma|\Phi|^2$ pairs of cherries contains $\gamma|\Phi|/4$ such pairs none of which share a cherry, then the latter case would imply that we applied $\gamma|\Phi|/8$ local improvements before.

In summary we showed that if for at least $\gamma|\mathcal{W}|/2 \geq \gamma \frac{\alpha t}{8k}$ (see (9)) many $K \in \mathcal{W}$ we can define a local improvement for $\gamma|\Phi|^2$ cherry pairs then we can aggregate local improvements leading to a β_{i+1} -hom(\mathcal{C}_ℓ)-tiling h'' with weight at least

$$w(h'') \geq w(h) + \min \left\{ \frac{\alpha}{2} \cdot \gamma \frac{\alpha t}{8k}, \frac{\beta_i}{4} \cdot \frac{\gamma}{8} |\Phi| \right\} \stackrel{(7),(8)}{\geq} w(h) + ct,$$

which would conclude the proof of Lemma 13.

Consequently, for the third step we may only consider those $K \in \mathcal{W}$ for which at most a γ -fraction of its cherry pairs $\mathcal{C}, \mathcal{C}'$ induce at least $4(k - \ell)$ edges or a 3-matching in $L_K(\mathcal{C}, \mathcal{C}')$. The degree condition (10) of $K \in \mathcal{W}$ translates to an average number of edges of at least $4(k - \ell) - 1 - 12\gamma(k - \ell)^2$ in the link graphs. So, as \mathcal{C}' was chosen big enough, at most $(\mathcal{C}' - 1)\gamma$ of the cherry pairs do not have exactly $4(k - \ell) - 1$ edges.

There can only be exactly $4(k - \ell) - 1$ edges and no 3-matching if there is a vertex on each side that has a complete neighbourhood, so these pairs are indeed extremal. Hence it remains to show that typically the special vertex $u \in \mathcal{C}$ in an extremal pair $L_K(\mathcal{C}, \mathcal{C}')$ is independent of K and \mathcal{C}' . So assume for a moment that there are $K, K' \in \mathcal{W}$ and two vertices u and v in some $\mathcal{C} \in \Phi$ such that u is a special vertex for an extremal pair $L_K(\mathcal{C}, \mathcal{C}')$ and v is special for some extremal pair $L_{K'}(\mathcal{C}, \mathcal{C}''')$. In this case we can define a local improvement by ‘‘splitting’’ the case with four edges above. Indeed choose four edges incident with u in $L_K(\mathcal{C}, \mathcal{C}')$ and four for v in $L_{K'}(\mathcal{C}, \mathcal{C}''')$. Assign weights $\frac{1}{2}\beta_i$ to the vertices of K and K' , and $\frac{k-2}{16(k-\ell-1)}\beta_i$ to the eight chosen edges. Set $h'(\mathcal{C}) = (1 - \frac{k-2}{4(k-\ell-1)})\beta_i$ and reduce the weights on \mathcal{C}' and \mathcal{C}'' by $\frac{k-2}{16(k-\ell-1)}\beta_i$ (or by $\frac{k-2}{8(k-\ell-1)}\beta_i$ in case $\mathcal{C}' = \mathcal{C}''$). Similarly as in the first step, this gives a local improvement of $\beta_i/4$.

If for at least $\gamma|\mathcal{W}|/2$ many $K \in \mathcal{W}$ we can define such a local improvement for $\gamma|\Phi|^2$ extremal pairs, we could aggregate them like in the second step, which would conclude the proof of Lemma 13. So for each cherry \mathcal{C} we may fix a vertex $u_{\mathcal{C}}$ having the properties of the claim. \square

We call $\mathcal{C} \in \Phi$ *good* if it is contained in at least $\frac{1}{2}|\Phi|$ extremal pairs for at least $\frac{1}{2}|\mathcal{W}|$ many $K \in \mathcal{W}$ and *bad* otherwise. It follows from Claim 14 that at most $5C'\gamma|\Phi|$ cherries are bad. Moreover, for every vertex $v \in V$ we denote by $\Phi_{\text{BAD}}(v)$ the number of bad cherries $\mathcal{C} \in \Phi$ that contain it.

To complete the proof of the lemma we will show that we find a matching M such that every vertex $v \in e \in M$ is contained in “many” good cherries. For each good cherry $\mathcal{C} \in \Phi$ there are a lot of choices for \mathcal{C}' and $K \in \mathcal{W}$ such that \mathcal{C} and \mathcal{C}' are an extremal pair for K . We will redistribute the weights to transfer weight from the non-special vertices of \mathcal{C} (and \mathcal{C}') to K , which will reduce the weight on v (since we will ensure that v is a non-special vertex). Repeating this for every $v \in e$ will allow us to obtain a local improvement for the tiling and repeating this for sufficiently many hyperedges $e \in M$ leads to the desired global improvement.

We define the function $a: V(\mathcal{R}) \rightarrow [0, 1]$ by $v \mapsto \beta_i \cdot \sum_{\mathcal{C} \in \Phi} \mathbf{1}_{\{v\}}(u_{\mathcal{C}})$, which assigns to a vertex the sum of weights used by special vertices. As any cherry contains $2(k - \ell)$ vertices, it is clear that $\sum_{v \in V(\mathcal{R})} a(v) \leq \frac{t}{2(k - \ell)}$ and, therefore, we can utilise the β -fractional non-extremality of \mathcal{R} for the weights $b(\cdot) = 1 - a(\cdot)$ and obtain

$$\sum_{e \in E(\mathcal{R})} \prod_{v \in e} b(v) \geq C\gamma \binom{t}{k}.$$

Since there are at most $5C'\gamma|\Phi|$ bad cherries, they contribute a weight of at most

$$\beta_i \sum_{v \in V(\mathcal{R})} |\Phi_{\text{BAD}}(v)| \leq \beta_i v(\mathcal{C}_\ell) \cdot 5C'\gamma|\Phi| \stackrel{(8)}{\leq} 5C'\gamma t. \quad (11)$$

We shall only use good cherries to redistribute weights for the desired β_{i+1} -hom-tiling, so we consider the function $b': V(\mathcal{R}) \rightarrow [0, 1]$ given by

$$b'(v) = \min \{0, b(v) - \beta_i \cdot |\Phi_{\text{BAD}}(v)|\}$$

and in view of (11) and $C' \ll C$ (cf. (7)) we have

$$\sum_{e \in E(\mathcal{R})} \prod_{v \in e} b'(v) \geq \frac{C}{2}\gamma \binom{t}{k}.$$

By a simple double counting argument there is a matching $M \subset E(\mathcal{R})$ with

$$\sum_{e \in M} \prod_{v \in e} b'(v) \geq \frac{C}{2}\gamma \cdot \frac{t}{k}$$

and since $b'(v) \in [0, 1]$ we have

$$\sum_{e \in M} k \cdot \min_{v \in e} \{b'(v)\} \geq \sum_{e \in M} k \prod_{v \in e} b'(v) \geq \frac{C}{2}\gamma t. \quad (12)$$

In particular, we may assume that $\min_{v \in e} \{b'(v)\} > 0$ for every $e \in M$, since this has no effect on inequality (12). Moreover, from the definition of the function $b'(\cdot)$ it then follows that $\min_{v \in e} \{b'(v)\} \geq \beta_i$ for every $e \in M$.

For each vertex $v \in \bigcup M$, we consider good cherries that contain v as a non-special vertex. Assume that we have $K \in \mathcal{W}$ and an extremal pair $\mathcal{C}, \mathcal{C}'$ such that v is a non-special vertex in \mathcal{C} . The link graph $L_K(\mathcal{C}, \mathcal{C}')$ contains all edges incident to the two special vertices. We define a *local weight shift* as follows. Assign weights $\frac{1}{2(k-\ell)-1} \cdot \frac{k-2}{4(k-\ell-1)}\beta_i$ to the vertices of all edges incident with exactly one of the special vertices, β_i to the vertices of K and set $h'(\mathcal{C}) = h'(\mathcal{C}') = (1 - \frac{k-2}{4(k-\ell-1)})\beta_i$. By similar calculations as before, this defines a valid β_{i+1} -hom-tiling h' with $w(h') = w(h)$. On the other hand, the weight of the vertex v and all other non-special vertices in $L_K(\mathcal{C}, \mathcal{C}')$ is reduced by $\frac{k-2}{4(k-\ell)-2}\beta_i$, i.e.

$$w_{h'}(v) = w_h(v) - \frac{k-2}{4(k-\ell)-2}\beta_i.$$

It follows from the definition of $b'(v)$ that we have at least $b'(v)/\beta_i$ many good cherries that contain v as non-special vertex and we shall apply at most $\min_{u \in e} \{b'(u)\}/\beta_i$ local weight shifts for a vertex $v \in e \in M$.

For every edge $e \in M$ we would like to apply these local weight shifts for every vertex $v \in e$, where we cycle through all k vertices and apply one shift at a time. In other words, we evenly reduce the weights on the vertices of e . Note that we can apply these local weight shifts using K, \mathcal{C} , and \mathcal{C}' unless we have saturated the vertices in K or used one of the cherries before. The procedure stops as soon as we reach a vertex for which no local weight shift is possible.

We first discuss the ideal case that this procedure does not stop, i.e. for every $e \in M$ and every $v \in e$ we applied $\min_{u \in e} \{b'(u)\}/\beta_i$ local weight shifts. In this case, for every $e \in M$ we reduced the weight of all vertices $v \in e$ by at least

$$\frac{1}{\beta_i} \min_{u \in e} \{b'(u)\} \cdot \frac{k-2}{4(k-\ell)-2}\beta_i = \frac{k-2}{4(k-\ell)-2} \min_{u \in e} \{b'(u)\}.$$

Consequently, we may appeal to Fact 12 increase the tiling on the edge e by the same amount. Repeating this for all $e \in M$, we obtain a β_{i+1} -hom-tiling h'' satisfying

$$w(h'') \geq w(h) + \sum_{e \in M} k \cdot \frac{k-2}{4(k-\ell)-2} \min_{u \in e} \{b'(u)\} \stackrel{(12)}{\geq} w(h) + \frac{C\gamma t}{2} \cdot \frac{k-2}{4(k-\ell)-2} \stackrel{(7)}{\geq} w(h) + ct,$$

which concludes the proof for this case.

In the case that the procedure stops, there is some $v \in e \in M$ and a good cherry \mathcal{C} for v such that \mathcal{C} cannot be used for a local weight shift for v . This means, since \mathcal{C} is a good cherry, that either there are $\frac{1}{2}|\mathcal{W}|$ many $K \in \mathcal{W}$ that contain a saturated vertex or that there are at least $\frac{1}{2}|\Phi|$ many cherries that were used in local weight shifts before. In the

case that $\frac{1}{2}|\mathcal{W}|$ many $K \in \mathcal{W}$ contain a saturated vertex, then each of these vertices was used in at least $\frac{\alpha}{2\beta_i}$ local weight shifts, so in total we have applied

$$\frac{1}{2}|\mathcal{W}| \cdot \frac{\alpha}{2\beta_i} \stackrel{(9)}{\geq} \frac{\alpha t}{8k} \cdot \frac{\alpha}{2\beta_i}$$

weight shifts. If on the other hand all $\frac{1}{2}|\Phi|$ possible cherries \mathcal{C}' were used in local weight shifts before and we have applied at least $\frac{1}{4}|\Phi|$ local weight shifts. As in the ideal case, using Fact 12, we conclude that we increase the tiling on the edges in M and obtain a β_{i+1} -hom-tiling h'' with

$$w(h'') \geq w(h) + \left(\min \left\{ \frac{\alpha^2 t}{16k\beta_i}, \frac{|\Phi|}{4} \right\} - k \right) \cdot \frac{(k-2)\beta_i}{4(k-\ell)-2} \stackrel{(7),(8)}{\geq} w(h) + ct,$$

which concludes the proof of Lemma 13. \square

Next we want to transfer the β -hom(\mathcal{C}_ℓ)-tiling of \mathcal{R} into a path-tiling of \mathcal{H} . For that purpose we will use the next lemma from [4].

Lemma 15. *Fix $k \geq 3$, $1 \leq \ell < k/2$ and $\varepsilon, d > 0$ such that $d > 2\varepsilon$. Let $m > \frac{k^2}{\varepsilon^2(d-\varepsilon)}$. Suppose $\mathcal{V} = (V_1, \dots, V_k)$ is an (ε, d) -regular k -tuple with*

$$|V_1| = \dots = |V_{2\ell}| = m \text{ and } |V_{2\ell+1}| = \dots = |V_k| = 2m.$$

Then there are at most $\frac{2k}{(d-\varepsilon)\varepsilon}$ vertex disjoint ℓ -paths that together cover all but at most $2k\varepsilon m$ vertices of \mathcal{V} . \square

Finally, by using Lemma 15 on the edges of the β -hom(\mathcal{C}_ℓ)-tiling of \mathcal{R} given by Lemma 13, we obtain a path-tiling from \mathcal{H} of the desired size.

Lemma 16 (Path-Tiling Lemma). *For all integers $k \geq 3$ and $1 \leq \ell < k/2$, there exist $C, \gamma_0 > 0$ such that for all $\alpha > 0$, $\gamma \leq \gamma_0$ there exists an integer s such that the following holds for all sufficiently large n . Let \mathcal{H} be a k -uniform hypergraph on n vertices and*

$$\delta_{k-2}(\mathcal{H}) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} - \gamma \right) \binom{n}{2}.$$

Then there is a family of at most s disjoint ℓ -paths that cover all but at most αn vertices of \mathcal{H} or \mathcal{H} is $(\ell, C\gamma)$ -extremal.

Proof. Let $k \geq 3$ and $1 \leq \ell < k/2$ be given. Let C' and γ'_0 be the constants given by Lemma 13 for k and ℓ . Set $C = 6C'$ and $\gamma_0 = \frac{\gamma'_0}{4}$, and let $\alpha > 0$ and $\gamma < \gamma_0$. Following the quantification of Lemma 13 with $\frac{\alpha}{2}$ and γ we obtain β and ε' and a sufficiently large t_0 . Let ε be sufficiently small. Then the weak regularity lemma (Lemma 9) for $\varepsilon_0 = \frac{\beta\varepsilon}{2} < \gamma^2$

and t_0 yields T_0 . Let s be a sufficiently large constant. Let \mathcal{H} be a k -uniform hypergraph on n vertices such that

$$\delta_{k-2}(\mathcal{H}) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} - \gamma \right) \binom{n}{2}$$

and n is sufficiently large. By the regularity lemma there exists an ε_0 -regular partition $V_0 \cup \dots \cup V_t$ of \mathcal{H} with $|V_1| = \dots = |V_t| = m$, $|V_0| \leq \varepsilon_0 n$ and $t_0 \leq t \leq T_0$ and the corresponding reduced hypergraph $\mathcal{R} = \mathcal{R}(\varepsilon_0, \gamma)$ on t vertices satisfies, by Lemma 10,

$$\deg_{\mathcal{R}}(K) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} - 4\gamma \right) \binom{t}{2}$$

for all but at most $\sqrt{\varepsilon_0} \binom{t}{k-2} \leq \varepsilon' \binom{t}{k-2}$ many $(k-2)$ -sets $K \in \binom{[t]}{k-2}$. We split the remainder of the proof in two cases, depending on whether \mathcal{R} is β -fractionally $(\ell, 4C'\gamma)$ -extremal.

Suppose that \mathcal{R} is not β -fractionally $(\ell, 4C'\gamma)$ -extremal. Then Lemma 13 implies that there exists a β -hom(\mathcal{C}_ℓ)-tiling h of \mathcal{R} with weight $(1 - \frac{\alpha}{2})t$. Let Φ^+ be the set of homomorphisms φ from \mathcal{C}_ℓ to \mathcal{R} with $h(\varphi) > 0$, which implies in fact $h(\varphi) \geq \beta$. We will use Lemma 15 to obtain ℓ -paths covering almost all vertices of \mathcal{H} and for this we split the partition classes according to the tiling h : let $\{R_1^\varphi, \dots, R_{2k-2\ell}^\varphi\}_{\varphi \in \Phi^+}$ be a family such that for all $\varphi \neq \varphi' \in \Phi^+$

- $R_i^\varphi \subset V_{\varphi(i)}$ for all $i \in [2k-2\ell]$,
- $R_i^\varphi \cap R_j^{\varphi'} = \emptyset$ for all $i, j \in [2k-2\ell]$,
- $|R_i^\varphi| = 2 \lfloor \frac{h(\varphi)m}{2} \rfloor$ for all $i \in [2k-2\ell]$.

For each $\varphi \in \Phi^+$ and all $i \in \{k-2\ell+1, \dots, k\}$ let $S_i^\varphi \cup U_i^\varphi = R_i^\varphi$ be a partition of R_i^φ into two classes of equal size. Note that

$$(R_1^\varphi, \dots, R_{k-2\ell}^\varphi, S_{k-2\ell+1}^\varphi, \dots, S_k^\varphi) \quad \text{and} \quad (U_{k-2\ell+1}^\varphi, \dots, U_k^\varphi, R_{k+1}^\varphi, \dots, R_{2k-2\ell}^\varphi)$$

are (ε, γ) -regular, where we used that $h(\varphi) \geq \beta$ for all $\varphi \in \Phi^+$. Then, with Lemma 15 we obtain at most $\frac{2k}{(\gamma-\varepsilon)\varepsilon}$ many ℓ -paths that cover all but $k\varepsilon|R_i^\varphi|$ vertices. Applying this to all homomorphisms $\varphi \in \Phi^+$ we obtain at most s many ℓ -paths.

We claim that the number of vertices in $V(\mathcal{H})$ that are not covered by these paths is less than αn . For this note that the uncovered vertices are the vertices from the partition class V_0 , the vertices that are not contained in any R_i^φ and those vertices in some R_i^φ that are not contained in any ℓ -path. At most $\frac{\alpha}{2}n$ vertices are not in any R_i^φ due to the weight of the β -hom(\mathcal{C}_ℓ)-tiling h and we lose at most $\frac{2t}{\beta}$ vertices due to the rounding in the definition of R_i^φ . The ℓ -paths cover all but a $(k\varepsilon)$ -fraction of vertices in $\bigcup_{i,\varphi} R_i^\varphi$. Consequently the total number of uncovered vertices is at most

$$\varepsilon_0 n + \frac{\alpha}{2}n + \frac{2t}{\beta} + k\varepsilon n < \alpha n.$$

Now suppose that \mathcal{R} is β -fractionally $(\ell, 4C'\gamma)$ -extremal. This means by definition that there is a function $b: V(\mathcal{R}) \rightarrow \{0\} \cup [\beta, 1]$ with

$$\sum_{v \in V(\mathcal{R})} b(v) \geq \frac{2(k-\ell)-1}{2(k-\ell)}t \quad \text{and} \quad \sum_{e \in E(\mathcal{R})} \prod_{v \in e} b(v) \leq 4C'\gamma \binom{t}{k}.$$

For each $i \in [t]$ we fix a subset $A_i \subseteq V_i$ with $|A_i| = \lfloor b(i)|V_i| \rfloor$ and define $B = \bigcup_{i \in [t]} A_i$. Thus, recalling the definition of the reduced hypergraph $\mathcal{R} = \mathcal{R}(\varepsilon_0, \gamma)$

$$\begin{aligned} e_{\mathcal{H}}(B) &\leq \sum_{e \in E(\mathcal{R})} \prod_{v \in e} \left(b(v) \frac{n}{t} \right) + \binom{t}{k} \gamma \left(\frac{n}{t} \right)^k + \varepsilon_0 \binom{t}{k} \left(\frac{n}{t} \right)^k + t \binom{n/t}{2} \binom{n}{k-2} \\ &\leq 4C'\gamma \binom{n}{k} + \gamma \binom{n}{k} + \varepsilon_0 \binom{n}{k} + \frac{k(k-1)}{2t} \binom{n}{k} \\ &\leq 5C'\gamma \binom{n}{k}. \end{aligned}$$

Note that

$$|B| \geq \left(\frac{2(k-\ell)-1}{2(k-\ell)}t \right) (1 - \varepsilon_0) \frac{n}{t} - t \geq \left(\frac{2(k-\ell)-1}{2(k-\ell)} - \varepsilon_0 \right) n.$$

Therefore, by adding at most $\varepsilon_0 n$ vertices from $V \setminus B$ to B we obtain a set B' with $|B'| = \left\lfloor \frac{2(k-\ell)-1}{2(k-\ell)}n \right\rfloor$ such that

$$e_{\mathcal{H}}(B') \leq e_{\mathcal{H}}(B) + \varepsilon_0 n \binom{n}{k-1} \leq 6C'\gamma \binom{n}{k} = C\gamma \binom{n}{k},$$

from which we conclude that \mathcal{H} is $(\ell, C\gamma)$ -extremal. \square

§3. PROOF OF THEOREM 4

Below we give the proof of the main technical result, which details the outline from Section 2.1 and is based on the lemmas from the last section.

Proof of Theorem 4. Let $0 < \xi < 1$ and let $k \geq 4$ and $1 \leq \ell < k/2$ be integers. Let C and γ_0 be given by the Path-Tiling Lemma (Lemma 16) for k and ℓ . Let $\gamma < \gamma_0$ be a sufficiently small constant. By the Absorbing Lemma (Lemma 7) for $\gamma, \zeta = \gamma, k$ and ℓ we obtain ε . Following the quantification of the Path-Tiling Lemma for $\alpha = \varepsilon/2$ and 3γ we obtain an integer s . We will use the Reservoir Lemma (Lemma 6) with $\eta = \frac{4(k-\ell)-1}{4(k-\ell)^2} - 2\gamma$, $\varepsilon' = \min\{\varepsilon/2, \gamma\}$, k , and $s+1$. Let $n \in (k-\ell)\mathbb{N}$ be sufficiently large and let \mathcal{H} be a k -uniform hypergraph on n vertices.

Suppose \mathcal{H} is not (ℓ, ξ) -extremal and

$$\delta_{k-2}(\mathcal{H}) \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} - \gamma \right) \binom{n}{2}.$$

Let \mathcal{A} be the absorbing path obtained with the Absorbing Lemma and let X_0 and Y_0 be the ends of \mathcal{A} . Then $|V(\mathcal{A})| \leq \gamma n$ and \mathcal{A} has the following absorption property: for every subset $U \subset V \setminus V(\mathcal{A})$ with $|U| \leq \varepsilon n$ and $|U| \in (k - \ell)\mathbb{N}$ there exists an ℓ -path $\mathcal{Q} \subset \mathcal{H}$ such that $V(\mathcal{Q}) = V(\mathcal{A}) \cup U$ and \mathcal{Q} has the ends X_0 and Y_0 .

Let $V' = (V \setminus V(\mathcal{A})) \cup \{X_0, Y_0\}$ and let $\mathcal{H}' = \mathcal{H}[V']$ be the subhypergraph of \mathcal{H} induced by V' . Note that

$$\delta_{k-2}(\mathcal{H}') \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} - 2\gamma \right) \binom{n}{2}.$$

The Reservoir Lemma guarantees the existence of a set $R \subset V'$ with $|R| \leq \varepsilon' n \leq \gamma n$ such that for every $j \leq s + 1$ and every family $(X_i, Y_i)_{i \in [j]}$ of mutually disjoint pairs of sets of ℓ vertices can be connected by paths that contain vertices of $\bigcup_{i \in [j]} (X_i \cup Y_i) \cup R$ only.

Let $V'' = V \setminus (V(\mathcal{A}) \cup R)$ and let $\mathcal{H}'' = \mathcal{H}[V'']$ be the subhypergraph of \mathcal{H} induced by V'' . Then

$$\delta_{k-2}(\mathcal{H}'') \geq \left(\frac{4(k-\ell)-1}{4(k-\ell)^2} - 3\gamma \right) \binom{n}{2}.$$

Now we apply the Path-Tiling Lemma to \mathcal{H}'' and obtain a family of at most s disjoint ℓ -paths that cover all but at most $\alpha|V''| \leq \alpha n$ vertices of \mathcal{H}'' , or \mathcal{H}'' is $(\ell, C\gamma)$ -extremal. Set $n'' = |V''|$ and suppose for a contradiction that \mathcal{H}'' is $(\ell, C\gamma)$ -extremal. Then there exists a set $B \subset V''$ such that $|B| = \lfloor \frac{2(k-\ell)-1}{2(k-\ell)} n'' \rfloor$ and $e(B) \leq C\gamma(n'')^k$. By adding at most $n - n'' \leq 2\gamma n$ vertices from $V \setminus B$ to B , we obtain a vertex set $B' \subset V$ such that $|B'| = \lfloor \frac{2(k-\ell)-1}{2(k-\ell)} n \rfloor$ and

$$e(B') \leq C\gamma(n'')^k + 2\gamma n \binom{n-1}{k-1} \leq \xi n^k,$$

a contradiction to the fact that \mathcal{H} is not (ℓ, ξ) -extremal. Therefore, we may assume that there exist disjoint ℓ -paths $\mathcal{P}_1, \dots, \mathcal{P}_j$ with $j \leq s$ that cover all but at most $\alpha|V''| \leq \alpha n$ vertices of \mathcal{H}'' .

For all $i \in [j]$, we denote the ends of \mathcal{P}_i by X_i and Y_i . Let $Y_{j+1} = Y_0$. By using the Reservoir Lemma to connect the family $(X_i, Y_{i+1})_{0 \leq i \leq j}$, we connect the ℓ -paths $\mathcal{A}, \mathcal{P}_1, \dots, \mathcal{P}_j$ to an ℓ -cycle $\mathcal{C} \subset \mathcal{H}$.

Let $U = V \setminus V(\mathcal{C})$ be the set of vertices not contained in \mathcal{C} , i.e. the vertices that were leftover in the reservoir R or uncovered by the path-tiling. We have $|U| \leq (\varepsilon' + \alpha)n \leq \varepsilon n$. Furthermore, since \mathcal{C} is an ℓ -cycle and $n \in (k - \ell)\mathbb{N}$, we have $|U| \in (k - \ell)\mathbb{N}$. Therefore, we can utilise the absorbing property of \mathcal{A} to replace \mathcal{A} in \mathcal{C} by a path \mathcal{Q} with the same ends as \mathcal{A} , obtaining a Hamiltonian ℓ -cycle of \mathcal{H} . \square

REFERENCES

- [1] J. de O. Bastos, G. O. Mota, M. Schacht, J. Schnitzer, and F. Schulenburg, *Loose Hamiltonian cycles forced by large $(k - 2)$ -degree – sharp version –*. In preparation. [↑1](#)
- [2] E. Buß, H. Hàn, and M. Schacht, *Minimum vertex degree conditions for loose Hamilton cycles in 3-uniform hypergraphs*, J. Combin. Theory Ser. B **103** (2013), no. 6, 658–678, DOI [10.1016/j.jctb.2013.07.004](#). MR3127586 [↑1](#), [2.2](#)
- [3] H. Hàn and M. Schacht, *Dirac-type results for loose Hamilton cycles in uniform hypergraphs*, J. Combin. Theory Ser. B **100** (2010), no. 3, 332–346, DOI [10.1016/j.jctb.2009.10.002](#). MR2595675 [↑1](#), [2.4](#)
- [4] J. Han and Y. Zhao, *Minimum codegree threshold for Hamilton ℓ -cycles in k -uniform hypergraphs*, J. Combin. Theory Ser. A **132** (2015), 194–223, DOI [10.1016/j.jcta.2015.01.004](#). MR3311344 [↑1](#), [1](#), [2.4](#)
- [5] ———, *Minimum vertex degree threshold for loose Hamilton cycles in 3-uniform hypergraphs*, J. Combin. Theory Ser. B **114** (2015), 70–96, DOI [10.1016/j.jctb.2015.03.007](#). MR3354291 [↑1](#), [1](#)
- [6] G. Y. Katona and H. A. Kierstead, *Hamiltonian chains in hypergraphs*, J. Graph Theory **30** (1999), no. 3, 205–212, DOI [10.1002/\(SICI\)1097-0118\(199903\)30:3<205::AID-JGT5>3.3.CO;2-F](#). MR1671170 [↑1](#)
- [7] P. Keevash, D. Kühn, R. Mycroft, and D. Osthus, *Loose Hamilton cycles in hypergraphs*, Discrete Math. **311** (2011), no. 7, 544–559, DOI [10.1016/j.disc.2010.11.013](#). MR2765622 [↑1](#)
- [8] D. König, *Graphok és matrixok*, Mat. Fiz. Lapok **38** (1931), 116–119 (Hungarian (German summary)). [↑2.4](#)
- [9] D. Kühn, R. Mycroft, and D. Osthus, *Hamilton ℓ -cycles in uniform hypergraphs*, J. Combin. Theory Ser. A **117** (2010), no. 7, 910–927, DOI [10.1016/j.jcta.2010.02.010](#). MR2652102 [↑1](#)
- [10] D. Kühn and D. Osthus, *Loose Hamilton cycles in 3-uniform hypergraphs of high minimum degree*, J. Combin. Theory Ser. B **96** (2006), no. 6, 767–821, DOI [10.1016/j.jctb.2006.02.004](#). MR2274077 [↑1](#)
- [11] V. Rödl, A. Ruciński, and E. Szemerédi, *A Dirac-type theorem for 3-uniform hypergraphs*, Combin. Probab. Comput. **15** (2006), no. 1-2, 229–251, DOI [10.1017/S0963548305007042](#). MR2195584 [↑1](#), [2.1](#), [2.3](#)
- [12] ———, *An approximate Dirac-type theorem for k -uniform hypergraphs*, Combinatorica **28** (2008), no. 2, 229–260, DOI [10.1007/s00493-008-2295-z](#). MR2399020 [↑1](#)
- [13] E. Szemerédi, *Regular partitions of graphs*, Problèmes combinatoires et théorie des graphes (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976), Colloq. Internat. CNRS, vol. 260, CNRS, Paris, 1978, pp. 399–401 (English, with French summary). MR540024 [↑2.4](#)

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, SÃO PAULO, BRAZIL
E-mail address: {josefran|mota}@ime.usp.br

FACHBEREICH MATHEMATIK, UNIVERSITÄT HAMBURG, HAMBURG, GERMANY
E-mail address: schacht@math.uni-hamburg.de
E-mail address: {jakob.schnitzer|fabian.schulenburg}@uni-hamburg.de